

# Lagrangian Floats for CBLAST

Eric A. D'Asaro

APL/UW 1013 NE 40<sup>th</sup> Str, Seattle, WA 98105

phone: (206) 685-2982 fax: (206) 543-6785 email: [dasaro@apl.washington.edu](mailto:dasaro@apl.washington.edu)

Award Number: N00014-00-1-0893

<http://opd.apl.washington.edu/~dasaro/HOME/>

## LONG-TERM GOALS

I seek to understand the dynamics of the ocean boundary layer beneath hurricanes and the resulting air-sea fluxes which drive it with the goal of improving ocean models at high wind speed.

## OBJECTIVES

To measure turbulence properties and fluxes in the ocean boundary layer beneath hurricanes and relate them to hurricane properties and fluxes measured by others. To model the measured boundary layer properties using Large Eddy Simulation (LES) techniques with the twin goals of testing the models and investigating the boundary layer physics using the models.

## APPROACH

**Measurements.** Neutrally buoyant Lagrangian floats were air-deployed into hurricanes during the 2002, 2003 and 2004 hurricane seasons. The floats are designed to be used in energetic turbulent flows such as those found in the top and bottom boundary layers of the ocean. A combination of accurate ballasting, compressibility matched to that of seawater and high drag is used to make these floats follow the motion of water parcels accurately (D'Asaro 2003). Water velocity is inferred from the motion of the floats; high frequency fluctuations in velocity can be used to infer dissipation rate (Lien and D'Asaro, 2006) and covariance of vertical velocity with scalars can be used to compute heat and other fluxes (D'Asaro, 2004).

**Modeling.** The LES modeling work is being conducted by Ramsey Harcourt and Eric D'Asaro. Our starting point is a standard LES scheme using a subgrid closure with active kinetic energy as implemented in Harcourt et. al (2002). The standard implementation of vortex force interaction between surface wave Stokes drift (Skylkingstad and Denbo, 1995; McWilliams *et al*, 1997) is modified to simulate the mean Lagrangian velocities measured by floats in the ocean. Careful attention has been paid to the role of surface waves in forcing boundary layer turbulence. This model includes the ability to simulate the trajectories of both perfectly Lagrangian and realistically imperfect floats.

## WORK COMPLETED

Work during FY08 focused on finishing the analysis of existing data and synthesizing the results over a wide range of wind speeds including hurricanes. It has been done jointly by Eric D'Asaro and Ramsey Harcourt. The work has focused on two questions:

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2008</b>		2. REPORT TYPE <b>Annual</b>		3. DATES COVERED <b>00-00-2008 to 00-00-2008</b>	
4. TITLE AND SUBTITLE <b>Lagrangian Floats For CBLAST</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>APL/UW,1013 NE 40th Str,Seattle,WA,98105</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>code 1 only</b>					
14. ABSTRACT <b>To measure turbulence properties and fluxes in the ocean boundary layer beneath hurricanes and relate them to hurricane properties and fluxes measured by others. To model the measured boundary layer properties using Large Eddy Simulation (LES) techniques with the twin goals of testing the models and investigating the boundary layer physics using the models.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES  <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			


- How well can Lagrangian floats measure ocean boundary layer turbulence? Although Lagrangian floats are excellent tools for measuring boundary layer processes, they are imperfect. The major cause of error is the buoyancy of the float, caused either by imperfect ballasting of the float or air bubbles attached to the float. This causes the float to spend more time near the surface and thus biases turbulence statistics measured from the float.
- What do LES models predict for turbulence levels in hurricanes? The prevailing paradigm for ocean boundary layer turbulence is that it is forced by both surface wind and waves. However, we do not have a quantitative understanding of the turbulence properties in such a boundary layer.

Both of these questions were addressed using LES modeling. An ensemble of LES model simulations spanning a wide range of oceanic conditions, including hurricanes, and including forcing by wind and surface waves via the Craik-Leibovich coupling. Lagrangian floats with realistic sizes, virtual masses, drag and a wide range of buoyancies were also simulated as in Harcourt, (2002). The wind stress and wave spectrum were forced to be consistent with the those of a pure wind sea. These simulation results were used to develop scaling laws for both the true properties of the ocean turbulence and the errors in float measurements of these properties. These are the two crucial components to rigorously assessing whether the float observations agree with the simulations. This final comparison will be completed under other funding.

## RESULTS

### How well can Lagrangian floats measure boundary layer turbulence?


Fig. 1 shows a typical result of many. Neutrally buoyant floats are mixed across the mixed layer by the energetic large turbulent eddies and rapidly become uniformly distributed in depth. Buoyant floats are not uniformly mixed, but rather spend more time near the surface. This biases the average vertical kinetic energy measured by floats high both because the kinetic energy is higher near the surface and because buoyant floats spend more time in the strongest-downward going plumes. Fig. 1 compares this bias, scaled by the measured vertical kinetic energy, to a simple measure of the depth distribution of the floats, the depth skewness. A good correlation exists. Since the skewness is an easily measured quantity, this result can be used to correct real observations. Under hurricane conditions, the skewness, and resulting bias, is typically quite small because the boundary layer turbulent velocities are very high. This adds strength to the previously reported good agreement between the float observations and model predictions in Hurricane Frances.



**Fig. 1. Bias in float measurement of boundary layer average vertical kinetic energy (vertical axis) as a function of the skewness of float depth (horizontal axis) for floats with upward buoyancies ranging from neutral to 80g both uniformly distributed in depth and surface intensified. Forcing characteristics vary from typical mid-latitude conditions (cases LW and MW) to hurricane conditions (cases H1 and H2). The depth skewness is a good predictor of the measurement error.**

## What do LES models predict for turbulence levels in hurricanes?

The same ensemble of LES models was used to find scaling laws for the turbulence levels in wind and wave forced ocean boundary layers. The most important result is shown in Fig. 2. The mixed layer average vertical kinetic energy is found to accurately scale with a Langmuir number, the square root of the ratio of surface wave Stokes drift to wind stress, in which change in Stokes drift from the top 25% of the boundary layer to the bottom of the layer is used instead of the more usual surface Stokes drift. These results are also been used to predict the boundary layer turbulence level as a function of wind speed and wave age. These results have been published in JPO.



**Fig. 2. Mixed layer average VKE from the ensemble of LES simulations scales accurately with the Surface Layer Langmuir number.**

## IMPACT/APPLICATIONS

The experimental and analysis techniques developed by this grant to study hurricanes continue to be used to study hurricanes under other funding and will be used in the upcoming ONR Typhoon Impact DRI.

## REFERENCES

- D'Asaro, Eric A. 2003 Performance of Autonomous Lagrangian Floats. *Journal of Atmospheric and Oceanic Technology* : Vol. 20, No. 6, pp. 896–911
- D'Asaro, E.A., (2004) Air-Sea Heat flux measurements from nearly neutrally buoyant floats, *J. Atmos. & Oceanic Tech.*, Vol. 21, No. 7, pp. 1086–1094
- Harcourt et al. (2001), Fully Lagrangian floats in Labrador Sea deep convection: Comparison of Numerical and experimental results, *J. Phys. Oceanogr.*, Vol. 32, No. 2, pp. 493-510.
- Large, W.G., J.C. McWilliams, and S.C. Doney, 1994: Oceanic vertical mixing: A review and a model with a nonlocal k-profile boundary layer parameterization, *Rev. Geophys.*, 32, 363-403.
- Lien, R-C. and E. D'Asaro and (2006), Measurement of turbulent kinetic dissipation rate using Lagrangian floats, *J. Atmos. & Oceanic Tech.*, 23, 964-976
- McWilliams, J. C., P. P. Sullivan and C.-H. Moeng. Langmuir turbulence in the ocean. *J. Fluid Mech.*, **334**, 1-30. 1997.

## PUBLICATIONS

- Harcourt, R, Eric D'Asaro, 2008, Large Eddy Simulation of Langmuir Turbulence in pure wind seas, *J. Phys. Oceanogr.*, **38**, 7 pp. 1542–1562, DOI: 10.1175/2007JPO3842.1 PUBLISHED